

A Modified Magnitude System that Produces Well-Behaved Magnitudes, Colors, and Errors Even for Low Signal-to-Noise Ratio Measurements

Robert H. Lupton^{1,2}, James E. Gunn², and Alexander S. Szalay³

ABSTRACT

We describe a modification of the usual definition of astronomical magnitudes, replacing the usual logarithm with an inverse hyperbolic sine function; we call these modified magnitudes ‘asinh magnitudes’. For objects detected at signal-to-noise ratios of greater than about five, our modified definition is essentially identical to the traditional one; for fainter objects (including those with a formally negative flux) our definition is well behaved, tending to a definite value with finite errors as the flux goes to zero.

This new definition is especially useful when considering the colors of faint objects, as the difference of two ‘asinh’ magnitudes measures the usual flux ratio for bright objects, while avoiding the problems caused by dividing two very uncertain values for faint objects.

The Sloan Digital Sky Survey (SDSS) data products will use this scheme to express all magnitudes in their catalogs.

Subject headings: catalogs, methods: statistical, techniques: photometric

1. Introduction

The advantages of using a logarithmic scale to measure astronomical fluxes are obvious: the magnitude scale is able to span a huge dynamic range, and when relative colors are needed they can be computed by simply differencing magnitudes measured in different bandpasses. These advantages are quite clear for bright objects where noise is not an issue.

¹For the SDSS collaboration.

²Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544

³Department of Physics and Astronomy, The Johns Hopkins University, Baltimore MD 21218

On the other hand, as fluxes become comparable to the sky and instrumental noise, the corresponding magnitudes are subject to large and asymmetric errors due to the singularity in the magnitude scale at zero flux; indeed, if a noisy measurement of an object’s flux happens to be negative, its magnitude is a complex number! Astronomers have generally handled these cases by specifying detection flags and somehow encoding the negative flux. These problems become even more pronounced as we work in multicolor space. An object can be well detected and measured in several of the bands, and still fail to provide a measurable flux in others. In flux space the object would be represented as a multivariate Gaussian probability distribution centered on its measured fluxes, not necessary all positive. Important information is lost by simply replacing a flux by an arbitrary 2- or 3- σ upper limit. In magnitude space, the error distribution of such an object has an infinite extent in some directions, making meaningful multicolor searches in a database impossible; all the non-detections in one or more bands must be isolated, and treated separately. One solution to this problem is to use linear units (e.g. Janskys), although this makes studies based on flux ratios (colors) inconvenient. This paper proposes an alternative solution, a modification of the definition of magnitudes, which preserves their advantages while avoiding their disadvantages.

2. The Inverse Hyperbolic Sine

We propose replacing the logarithm in the traditional definition of a magnitude with an inverse hyperbolic sine function. This function becomes a logarithm for large values of its argument, but is linear close to the origin.

$$\sinh^{-1}(x) = \ln \left[x + \sqrt{x^2 + 1} \right] \rightarrow \begin{cases} \operatorname{sgn}(x) \ln |2x|, & \text{if } |x| \gg 1 \\ x, & \text{if } |x| \lesssim 1 \end{cases} \quad (1)$$

The usual apparent magnitude m can be written in terms of the dimensionless normalized flux $x \equiv f/f_0$ as

$$m \equiv -2.5 \log_{10} x = -(2.5 \log_{10} e) \ln x \equiv -a \ln x. \quad (2)$$

where f_0 is the flux of an object with magnitude 0.0, and $a \equiv 2.5 \log_{10} e = 1.08574$ is Pogson’s ratio (Pogson 1856).

Let us define the new magnitude μ as

$$\mu(x) \equiv -a \left[\sinh^{-1} \left(\frac{x}{2b} \right) + \ln b \right] \quad (3)$$

Here a and b are constants, and b is an arbitrary ‘softening’ which determines the flux level at which linear behaviour sets in. After some discussion, we have adopted the name ‘asinh

magnitudes’ for μ . Consider the asymptotic behaviour of μ , for both high and low x :

$$\lim_{x \rightarrow \infty} \mu(x) = -a \ln x = m \qquad \lim_{x \rightarrow 0} \mu(x) = -a \left[\frac{x}{2b} + \ln b \right]. \quad (4)$$

Thus for $x \rightarrow \infty$, μ approaches m , for any choice of b . On the other hand when $|x| \lesssim b$, μ is linear in x ; for $x \ll -b$, we gradually return to logarithmic behaviour, although this regime is never of astronomical interest for any reasonable choice of b . Intuition suggests that b should be chosen to be comparable to the (normalized) flux of an object with a signal-to-noise ratio of about one; the following section discusses the choice of b , and the related question of μ ’s error distribution.

3. The Errors in μ and the choice of the Softening Parameter b

Although the softening parameter b can be any positive number, we show below that one particular selection has a couple of attractive features. In making our choice, we use the following guiding principles: 1) Since asinh magnitudes are being introduced to avoid the problems that classical magnitudes manifest in low signal-to-noise data, the differences between asinh and classical magnitudes should be minimized for high signal-to-noise data. 2) We should minimize μ ’s variance at low flux levels. The latter is not strictly required as a too-small value of b merely stretches out μ ’s scale along with its errors, but it is convenient if μ ’s variance at zero flux is comparable to its variance at a signal-to-noise ratio of a few.

In reality, our measurement of the normalized flux x will be noisy, with variance σ^2 . We wish to choose b to minimize μ ’s variance, while keeping the difference $m - \mu$ small. Let us therefore compute the variances of m and μ (keeping only the linear terms in their Taylor series), and also their difference; arrows indicate the asymptotic behavior as $x \rightarrow 0$:

$$\begin{aligned} \text{Var}(m) &= \frac{a^2 \sigma^2}{x^2} && \rightarrow \frac{a^2 \sigma^2}{x^2} \\ \text{Var}(\mu) &= \frac{a^2 \sigma^2}{4b^2 + x^2} && \rightarrow \frac{a^2 \sigma^2}{4b^2} \\ m - \mu &= a \ln \left[\frac{1 + \sqrt{1 + 4b^2/x^2}}{2} \right] && \rightarrow -a \operatorname{sgn}(x) \ln \left(\frac{|x|}{b} \right) \end{aligned} \quad (5)$$

What are the disadvantages of taking either too low or too high a value for b ? Choosing a low value of b causes the difference $m - \mu$ to become smaller, i.e. the two magnitudes track each other better, but unfortunately μ ’s variance at $x = 0$ varies as $1/b^2$. Choosing too high a value has the opposite effect: the difference explodes at low values of x , simply due to the

singularity in the logarithm in the definition of m . At the same time, μ 's variance remains small.

In order to balance these two competing effects, we shall determine b by minimizing a penalty function containing terms due to both, added in quadrature.

The difference between the two magnitudes, normalised by m 's standard deviation, is given by

$$\delta(x) = \frac{m(x) - \mu(x)}{\sqrt{\text{Var}(m)}} \equiv \frac{b}{\sigma} F\left(\frac{x}{b}\right), \quad \text{where} \quad F(y) = y \ln \left[\frac{1 + \sqrt{1 + 4/y^2}}{2} \right]. \quad (6)$$

The function $F(y)$ has a maximum value of approximately 0.5 at $y = 0.7624$, so the largest possible deviation between the two magnitude scales is

$$(m - \mu)_{\text{max}} \approx \frac{b}{2\sigma} \sqrt{\text{Var}(m)} \quad (7)$$

The other ‘cost’ associated with the choice of b is the size of the error box for μ at $x = 0$, which is

$$\sqrt{\text{Var}(\mu)|_{x=0}} = \frac{a\sigma}{2b} \quad (8)$$

The total penalty can be obtained by adding these two costs in quadrature:

$$\epsilon = \delta_{\text{max}}^2 + \text{Var}(\mu)|_{x=0} = \frac{b^2}{4\sigma^2} + \frac{a^2\sigma^2}{4b^2} = \frac{a}{4} \left[\frac{b^2}{a\sigma^2} + \frac{a\sigma^2}{b^2} \right] \quad (9)$$

which has the obvious minimum at $b^2 = a\sigma^2$. Thus the optimal setting is the value $b = \sqrt{a}\sigma = 1.042\sigma$. As expected, b is approximately equal to the noise in the flux. This choice of b leads to $m - \mu$ having a maximum value of $0.52\sqrt{\text{Var}(m)}$, implying that the difference between the two magnitudes is always smaller than the uncertainty in m (see figure 1). If the error in its measured flux is 1σ , the error in μ is ± 0.52 . If the flux errors are Gaussian, so, to leading order, are the errors in μ as the transformation from counts to μ is linear for $|x| \lesssim b$.

Figure 2 shows m and μ as a function of signal-to-noise ratio, for this choice of b , along with their 1σ errors.

4. Application to Real Data

We have been working in terms of $x \equiv f/f_0$, but it is usually more convenient to use the measured fluxes directly; In terms of the *non*-normalised flux f , the expressions for m ,

μ , and $\text{Var}(\mu)$ become

$$m = m_0 - 2.5 \log_{10} f, \quad (10)$$

$$\mu = (m_0 - 2.5 \log_{10} b') - a \sinh^{-1}(f/2b') \quad (11)$$

and

$$\text{Var}(\mu) = \frac{a^2 \sigma'^2}{4b'^2 + f^2} \approx \frac{a^2 \sigma'^2}{4b'^2} \quad (12)$$

where $m_0 \equiv 2.5 \log_{10}(f_0)$, $b' \equiv f_0 b$, and $\sigma' \equiv f_0 \sigma$ are measured in real flux units (e.g. counts).

An object with no measured flux in a given band has a μ value of $m_0 - 2.5 \log_{10} b'$ (equation 11), in other words the classical magnitude of an object with a flux of b' . We note in passing that this value $\mu(0)$ is a convenient measure of the depth of a survey, containing information about both the noise properties of the sky and the image quality.

In the discussion above we considered the idealized case of all objects having the same error, dominated by sky noise. This case covers most objects found in a given deep survey, as most are detected at the flux limit and are typically at most marginally resolved. For bright objects, of course, the difference between m and μ is entirely negligible.

The optimal measure of the flux of a faint stellar object is given by convolving its image with the PSF. If the noise is dominated by the sky and detector, the variance of the measured flux is independent of the object's brightness, and is given by the background variance per unit area multiplied by the effective area of the PSF ($4\pi\alpha^2$ if the PSF is Gaussian with FWHM $2\sqrt{2 \ln 2} \alpha$). If we decide upon a typical seeing quality and sky brightness for a given band, this defines σ 's nominal value, σ_0 , which sets b once and for all. Each band has its own value of b .

As observing conditions change so do measurement errors, with the result that the error in μ for very faint objects is no longer exactly the 0.52 magnitudes that it would be under canonical conditions. Whenever a precise error is needed for a given object's μ , it may be found by converting μ back to flux, or by applying equation 12; for faint objects this reduces to multiplying the quoted error by σ/σ_0 . Failure to apply such a correction would mean that the quoted errors on μ were wrong.

It would be possible to choose b separately for different parts of the sky, but this would make the conversion of μ back to flux impractically complicated, and a significant source of mistakes for users of the data. The behaviour of μ as b changes is reasonably benign; the error at zero flux varies only linearly with b , as does the flux where m and μ begin to diverge.

Care is also required whenever the measured flux has different noise properties, for

example if the flux is measured within a circular aperture or a given isophote rather than using a PSF. In this case, the appropriate value of σ may be much larger from the one used to set b , with the consequence that the error in μ at zero flux considerably underestimates the true uncertainty (the other case, where the effective aperture is smaller than the PSF, is unlikely to occur in practice). It would, of course, be possible if confusing to choose a different set of b values for each type of (fixed size) aperture, although it seems unlikely that this would really be a good idea. As the discussion in section 3 showed the consequences of even a grossly incorrect value of b are not catastrophic; the asinh magnitudes still reduce to our familiar magnitudes for bright objects, and are still well defined for negative values.

One place where special care will be needed is in measures of surface brightness, where m and μ can depart quite strongly from one another even at levels where the flux is well determined. It may prove desirable to use a different value of b for such measurements; they are after all never directly compared with total magnitudes.

Fan et al. (1999a) have used the asinh magnitude system to search for high- z quasars in preliminary SDSS data; examples of color-color plots employing asinh magnitudes may be found in Fan et al. (1999b).

5. Asinh Magnitudes and Colors

The ratio of two low signal-to-noise ratio measurements (for example, an object’s color) is statistically badly behaved (indeed, for Gaussian distributions if the denominator has zero mean, the ratio follows a Cauchy distribution and accordingly has no mean, let alone a variance!). What is the behavior of our asinh magnitudes when used to measure colors?

For objects detected at high signal-to-noise ratio, the difference in μ measured in two bands is simply a measure of the relative flux in the two bands. For faint objects this is no longer true, although the difference is well behaved. A non-detection in two bands has a well defined ‘color’ ($\mu_1(0) - \mu_2(0)$). As discussed above, the error in this color is approximately $0.75\sigma/\sigma_0$ magnitudes, assuming independent errors in the two bands. Equivalently, such a non-detection can be represented by an ellipsoid in multi-color space, centered at the point corresponding to zero flux in all bands, with principal axes $0.52\sigma/\sigma_0$ (in general σ/σ_0 will be different in each band).

As an illustration of the instability of the traditional definition of color for faint objects, consider two objects that have almost identical colors but which are near the detection limit of a survey. Their asinh colors will be very similar, but (due to the singularity of the logarithm as the flux goes to zero) their classical magnitudes may differ by an arbitrarily

large amount.

Figure 3 shows the results of a simple Monte-Carlo simulation. We took a set of ‘objects’ with 2.51 times as much flux in one band in the other (one magnitude), added Gaussian noise of fixed variance to each measurement, and tabulated the color measured using both classical and asinh magnitudes. The left hand panel shows the flux ratio, $\Delta m \equiv m_1 - m_2$, and $\Delta \mu \equiv \mu_1 - \mu_2$ as a function of signal-to-noise ratio; the right hand panels show histograms of their distribution in the range $1 \leq S/N \leq 3$. At the right side of the plot, where the noise is less important, both $\Delta m \equiv m_1 - m_2$ and $\Delta \mu \equiv \mu_1 - \mu_2$ tend to -1, the correct value. As the noise becomes more important, the errors on the Δm plot grow (and an increasing fraction of points in the left panel is simply omitted as their fluxes are zero or negative). The $\Delta \mu$ plot shows the ‘color’ tending to its value at zero flux, in this case 0.0, as the signal-to-noise ratio drops.

6. Summary

We have shown that an innovative use of inverse hyperbolic sines for a new magnitude scale can overcome most deficiencies of traditional magnitudes, while preserving their desirable features. The defining equations are

$$\mu = (m_0 - 2.5 \log_{10} b') - a \sinh^{-1}(f/2b') \equiv \mu(0) - a \sinh^{-1}(f/2b')$$

and

$$\text{Var}(\mu) = \frac{a^2 \sigma'^2}{4b'^2 + f^2} \approx \frac{a^2 \sigma'^2}{4b'^2}$$

where $a \equiv 2.5 \log_{10} e$, f is the measured flux, σ' the error in f due to the sky and detector, and b' a softening parameter (Equations 11 and 12).

The principal advantages of these ‘asinh’ magnitudes are their equivalence to classical magnitudes when errors are negligible, their ability to represent formally negative fluxes, and their well behaved error distribution as the measured flux goes to zero. For high signal-to-noise ratios the difference of two asinh magnitudes is a measure of the flux ratio, while for noisy detections it becomes the statistically preferable flux difference; this allows meaningful color cuts even when an object is barely detected in some bands. Additionally, $\mu(0)$ provides a convenient way of summarizing the photometric depth of a survey for point-like objects, containing information about both the noise level of the system and the image quality.

Asinh magnitudes will be used in the SDSS catalogs.

The authors would like to thank Xiaohui Fan, Don Schneider, and Michael Strauss for

helpful comments on various versions of this paper, and Gillian Knapp for demonstrating that the whole question was more subtle than we had realised.

The Sloan Digital Sky Survey (SDSS) is a joint project of The University of Chicago, Fermilab, the Institute for Advanced Study, the Japan Participation Group, The Johns Hopkins University, Princeton University, the United States Naval Observatory, and the University of Washington. Apache Point Observatory, site of the SDSS, is operated by the Astrophysical Research Consortium. Funding for the project has been provided by the Alfred P. Sloan Foundation, the SDSS member institutions, the National Aeronautics and Space Administration, the National Science Foundation and the U.S. Department of Energy.

REFERENCES

- Fan X. et al., 1999a, AJ, submitted.
Fan X. et al., 1999b, in preparation.
Pogson N.R., MNRAS **17**, 12, 1856.




Fig. 1.— The behavior of $m - \mu$ as a function of signal-to-noise ratio, x/σ . The solid line is the value of $m - \mu$ and the region between the dotted lines corresponds to the $\pm 1\sigma$ error region for m . The dashed lines are drawn at ± 0.01 .




Fig. 2.— The behavior of m and μ , and their respective errors, as a function of signal-to-noise ratio, x/σ . The solid line is the value of μ and the region between the dotted lines its $\pm 1\sigma$ error region; the points with errorbars are the classical magnitudes, m . We have arbitrarily chosen a zeropoint of $\mu = 25.0$ for an object with no flux. One other feature of our modified magnitudes is apparent from this figure, namely that the error band on μ is nearly symmetrical, while the errors in m are strongly skewed at faint magnitudes. For S/N ratios of less than about two, $m - \mu$ exceeds the value $0.52\sqrt{\text{Var}(m)}$ quoted in the main body of the paper; this is due to the breakdown of the linear approximation used to calculate m 's variance.




Fig. 3.— The results of a Monte-Carlo simulation of the colors of a set of objects. The bottom panel shows the ratio of the brighter to the fainter measurement; it is clear that many of the low signal-to-noise points have negative flux ratios. The center panel shows the same points, but now in terms of the magnitude difference of the two detections. We have simply omitted points for which one or both fluxes were negative, although they do appear in the histogram. Finally, the top panel shows the difference in the asinh magnitudes. The ‘color’ at zero flux was taken to be 0.0 for this simulation.